

Inula aschersoniana

Habitus and growth type

Life span: **Perennial**

Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Seed mass [mg]: **0.59**

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Ellenberg-type indicator values

Light indicator value: **8.4**

Temperature indicator value: **7.8**

Moisture indicator value: **3.3**

Reaction indicator value: **7.4**

Nutrient indicator value: **3.1**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.91**

Disturbance frequency (herb layer): **1.81**

Disturbance severity: **0.42**

Disturbance severity (herb layer): **0.35**

Mowing frequency: **0.1**

Grazing pressure: **0.31**

Soil disturbance: **0.18**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [KC \(ASP\) *Asplenietea trichomanis*](#)

Broad habitat

Occurrence in broad habitats: **Sparsely vegetated (incl. rock and scree)**