Inula ensifolia

Habitus and growth type

Plant height [m]: **0.28**
Life span: **Perennial**
Life form: **Hemicryptophyte**

Flower

Flowering period: **June-August**

Fruit, seed and dispersal

Seed mass [mg]: **0.52**
Dispersal mode: **Anemochory**
Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.1**
Temperature indicator value: **6.7**
Moisture indicator value: **3.3**
Reaction indicator value: **8**
Nutrient indicator value: **2.3**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.77**
Disturbance frequency (herb layer): **1.67**
Disturbance severity: **0.47**
Disturbance severity (herb layer): **0.37**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Mowing frequency: **0.3**
Grazing pressure: **0.3**
Soil disturbance: **0.14**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CL (GER) *Trifolio-Geranietea sanguinei*, DA (FES) *Festuco-Brometea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **R16 Perennial rocky grassland of Central and South-Eastern Europe**

Constant species of EUNIS habitats: **R16 Perennial rocky grassland of Central and South-Eastern Europe**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Forest, Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz