

Jovibarba heuffelii

Habitus and growth type

Life span: **Perennial**

Life form: **Chamaephyte**

Fruit, seed and dispersal

Dispersal mode: **Myrmecochory**

Dispersal distance class: **3**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Disturbance indicator values

Disturbance frequency: **0.75**

Disturbance frequency (herb layer): **1.25**

Disturbance severity: **0.36**

Disturbance severity (herb layer): **0.27**

Mowing frequency: **0.06**

Grazing pressure: **0.24**

Soil disturbance: **0.14**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CK \(SED\) Sedo-Scleranthetea](#), [KC \(ASP\) Asplenietea trichomanis](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [U32 Temperate high-mountain siliceous inland cliff](#)

Constant species of EUNIS habitats: [U32 Temperate high-mountain siliceous inland cliff](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Sparsely vegetated (incl. rock and scree)**