Juncus capitatus

Habitus and growth type

Plant height [m]: **0.08**
Life span: **Annual**
Life form: **Therophyte**

Flower

Flowering period: **March-May**

Fruit, seed and dispersal

Seed mass [mg]: **0.01**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**
Substrate reaction relationship: **Acidic**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.1**
Nutrient indicator value: **2**
Salinity indicator value: **0.4**

Disturbance indicator values

Disturbance frequency: **1.92**
Disturbance frequency (herb layer): **2.06**
Disturbance severity: **0.51**
Disturbance severity (herb layer): **0.51**
Mowing frequency: **0.2**
Grazing pressure: **0.23**
Soil disturbance: **0.31**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **OC (ISO) *Isoëto-Nanojuncetea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **Q62 Periodically exposed shore with stable, mesotrophic sediments with pioneer or ephemeral vegetation**, **R1R Mediterranean to Atlantic open, dry, acid and neutral grassland**

Broad habitat

Occurrence in broad habitats: **Coastal saltmarsh, Coastal beach, dune or shingle, Spring, Wetland, Grassland (non-alpine, non-saline), Saline vegetation, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **4**
Continentality amplitude: **4**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz