Kickxia spuria

Habitus and growth type

Plant height [m]: **0.16**
Life span: **Annual**
Life form: **Therophyte**

Leaf

Specific leaf area [mm^2/mg]: **23.72**

Flower

Flowering period: **May-September**

Fruit, seed and dispersal

Seed mass [mg]: **0.36**
Dispersal mode: **Local non-specific dispersal, Anthropochory**
Dispersal distance class: **1, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.4**
Moisture indicator value: **4.2**
Reaction indicator value: **6.8**
Salinity indicator value: **0.1**

Disturbance indicator values

Disturbance frequency: **2.01**
Disturbance frequency (herb layer): **2.02**
Disturbance severity: **0.83**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance severity (herb layer): **0.83**
Mowing frequency: **0.64**
Grazing pressure: **0.17**
Soil disturbance: **0.75**

Habitat and sociology

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Synanthropic**

Distribution

Continentality: **4**
Continentality amplitude: **4**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz