

Koeleria pyramidata

Habitus and growth type

Plant height [m]: **0.52**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **9.11**

Flower

Flowering period: **May-August**

Fruit, seed and dispersal

Seed mass [mg]: **1.71**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.6**

Temperature indicator value: **5.7**

Moisture indicator value: **3.8**

Reaction indicator value: **7**

Nutrient indicator value: **2.4**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.32**

Disturbance frequency (herb layer): **1.84**

Disturbance severity: **0.49**

Disturbance severity (herb layer): **0.47**

Mowing frequency: **0.82**

Grazing pressure: **0.3**

Soil disturbance: **0.15**

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: [R1A Semi-dry perennial calcareous grassland \(meadow steppe\)](#), [R1C Desert steppe](#)

Constant species of EUNIS habitats: [R15 Continental dry rocky steppic grassland and dwarf scrub on chalk outcrops](#), [R1A Semi-dry perennial calcareous grassland \(meadow steppe\)](#), [R1B Continental dry grassland \(true steppe\)](#), [R1C Desert steppe](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Oromediterranean scrub, Scrub, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality: **6**

Continentality amplitude: **5**