

# *Lactuca plumieri*

## Habitus and growth type

Plant height [m]: **0.68**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Flower

Flowering period: **May-July**

## Fruit, seed and dispersal

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **5.7**

Temperature indicator value: **3.8**

Reaction indicator value: **5.3**

Nutrient indicator value: **6.3**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.28**

Disturbance frequency (herb layer): **1.39**

Disturbance severity: **0.56**

Disturbance severity (herb layer): **0.23**

Mowing frequency: **0.06**

Grazing pressure: **0.18**

Soil disturbance: **0.13**

## **Habitat and sociology**

Syntaxon

Diagnostic species of phytosociological classes: [CS \(MUL\) \*Mulgedio-Aconitetea\*](#)

Broad habitat

Occurrence in broad habitats: **Wetland, Alpine-subalpine and arctic grassland, Scrub, Forest**

## **Distribution**

Continentality: **2**

Continentality amplitude: **2**