

Larix sibirica

Habitus and growth type

Plant height [m]:

Life span: **Perennial**

Life form: **Phanerophyte, Tree**

Fruit, seed and dispersal

Seed mass [mg]: **9.7**

Dispersal mode: **Anemochory**

Dispersal distance class: **4**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Ellenberg-type indicator values

Moisture indicator value: **4**

Disturbance indicator values

Disturbance frequency: **0.1**

Disturbance frequency (herb layer): **1.25**

Disturbance severity: **0.7**

Disturbance severity (herb layer): **0.11**

Mowing frequency: **0**

Grazing pressure: **0.18**

Soil disturbance: **0.14**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [BA \(PIC\) *Vaccinio-Piceetea*](#), [BC \(BRA\) *Brachypodio pinnati-Betuletea pendulae*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [T3H *Larix* light taiga](#)

Constant species of EUNIS habitats: [T3H *Larix* light taiga](#)

Dominant species of EUNIS habitats: [T3H Larix light taiga](#)

Broad habitat

Occurrence in broad habitats: **Forest**