

Lathyrus bauhini

Habitus and growth type

Plant height [m]: **0.35**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **5**

Temperature indicator value: **6.7**

Moisture indicator value: **5**

Reaction indicator value: **8.5**

Nutrient indicator value: **2.5**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.05**

Disturbance frequency (herb layer): **1.34**

Disturbance severity: **0.69**

Disturbance severity (herb layer): **0.13**

Mowing frequency: **0.01**

Grazing pressure: **0.24**

Soil disturbance: **0.1**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CL \(GER\) *Trifolio-Geranietea sanguinei*](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**

Distribution

Continentality: **5**

Continentality amplitude: **1**