

Lathyrus clymenum

Habitus and growth type

Plant height [m]: **0.75**

Life span: **Annual**

Life form: **Therophyte**

Flower

Flowering period: **March-June**

Fruit, seed and dispersal

Seed mass [mg]:

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.2**

Temperature indicator value: **9.3**

Moisture indicator value: **3.7**

Reaction indicator value: **6.3**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.51**

Disturbance frequency (herb layer): **1.49**

Disturbance severity: **0.74**

Disturbance severity (herb layer): **0.53**

Mowing frequency: **0.29**

Grazing pressure: **0.32**

Soil disturbance: **0.22**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [**FF \(LYG\) *Lygeo sparti-Stipetea tenacissimae***](#), [**QC \(CHE\) *Chenopodietea***](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Synanthropic**