Lathyrus clymenum

Habitus and growth type

Plant height [m]: **0.75**
Life span: **Annual**
Life form: **Therophyte**

Flower

Flowering period: **March-June**

Fruit, seed and dispersal

Seed mass [mg]:
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **symbiosis with rhizobia**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.2**
Temperature indicator value: **9.3**
Moisture indicator value: **3.7**
Reaction indicator value: **6.3**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.51**
Disturbance frequency (herb layer): **1.49**
Disturbance severity: **0.74**
Disturbance severity (herb layer): **0.53**
Mowing frequency: **0.29**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Grazing pressure: **0.32**
Soil disturbance: **0.22**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **FF (LYG) *Lygeo sparti-Stipetea tenacissimae*, QC (CHE) *Chenopodietea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **R1E Mediterranean tall perennial dry grassland**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Synanthropic**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz