Lathyrus niger

Habitus and growth type

Plant height [m]: **0.47**
Life span: **Perennial**
Life form: **Hemicryptophyte, Geophyte**

Flower

Flowering period: **April-June**

Fruit, seed and dispersal

Seed mass [mg]: **17.54**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **symbiosis with rhizobia**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Temperature indicator value: **6.4**
Moisture indicator value: **3.8**
Reaction indicator value: **6.7**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.03**
Disturbance frequency (herb layer): **1.31**
Disturbance severity: **0.69**
Disturbance severity (herb layer): **0.12**
Mowing frequency: **0.02**
Grazing pressure: **0.2**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Soil disturbance: **0.1**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CA (FAG) *Carpino-Fagetea sylvaticae*, CB (PUB) *Quercetea pubescentis***

EUNIS habitat

Diagnostic species of EUNIS habitats: **T19 Temperate and submediterranean thermophilous deciduous forest, T1E Carpinus and Quercus mesic deciduous forest**

Constant species of EUNIS habitats: **T19 Temperate and submediterranean thermophilous deciduous forest**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest**

Distribution

Continentality: **5**

Continentality amplitude: **3**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz