

# *Lathyrus venetus*

## Habitus and growth type

Plant height [m]: **0.34**

Life span: **Perennial**

Life form: **Hemicryptophyte, Geophyte**

## Flower

Flowering period: **March-May**

## Fruit, seed and dispersal

Seed mass [mg]: **16.29**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **4.3**

Temperature indicator value: **6.9**

Moisture indicator value: **5.1**

Reaction indicator value: **6.7**

Nutrient indicator value: **5**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.01**

Disturbance frequency (herb layer): **1.3**

Disturbance severity: **0.7**

Disturbance severity (herb layer): **0.11**

Mowing frequency: **0**

Grazing pressure: **0.2**

Soil disturbance: **0.1**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [CB \(PUB\) Quercetea pubescentis](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [T1G Alnus cordata forest](#)

Constant species of EUNIS habitats: [T1G Alnus cordata forest](#), [T33 Mediterranean mountain Abies forest](#)

### Broad habitat

Occurrence in broad habitats: **Forest**