

Legousia speculum-veneris

Habitus and growth type

Plant height [m]: **0.21**

Life span: **Annual**

Life form: **Therophyte**

Leaf

Specific leaf area [mm²/mg]: **65.57**

Flower

Flowering period: **March-July**

Fruit, seed and dispersal

Seed mass [mg]: **0.18**

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **1, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.6**

Temperature indicator value: **7.7**

Moisture indicator value: **4.2**

Reaction indicator value: **7.6**

Nutrient indicator value: **4.2**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.66**

Disturbance frequency (herb layer): **2.01**

Disturbance severity: **0.83**

Disturbance severity (herb layer): **0.82**

Mowing frequency: **0.57**

Grazing pressure: **0.16**

Soil disturbance: **0.78**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [QC \(CHE\) *Chenopodietea*](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Sparsely vegetated (incl. rock and scree), Synanthropic**

Distribution

Continentality: **5**

Continentality amplitude: **4**