Lemna minor

Habitus and growth type

Plant height [m]: **0.01**
Life span: **Perennial**
Life form: **Hydrophyte**

Leaf

Specific leaf area [mm^2/mg]: **18.7**

Flower

Flowering period: **April-October**

Fruit, seed and dispersal

Seed mass [mg]:
Dispersal mode: **Epizoochory**
Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Water**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Slightly saline or brackish**

Ellenberg-type indicator values

Light indicator value: **7**
Moisture indicator value: **11.1**
Nutrient indicator value: **5.6**
Salinity indicator value: **0.6**

Disturbance indicator values

Disturbance frequency: **1.38**
Disturbance frequency (herb layer): **1.65**
Disturbance severity: **0.27**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance severity (herb layer): **0.26**
Mowing frequency: **0.16**
Grazing pressure: **0.05**
Soil disturbance: **0.07**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **NA (LEM) *Lemnetea***

EUNIS habitat

Diagnostic species of EUNIS habitats: **P3b Fresh-water small pleustophyte vegetation**, **P3c Fresh-water large pleustophyte vegetation**, **P3d Fresh-water submerged vegetation**, **P3e Fresh-water nymphaeid vegetation**, **V14 Inundated or inundatable cropland, including rice fields**

Constant species of EUNIS habitats: **P3b Fresh-water small pleustophyte vegetation**, **P3c Fresh-water large pleustophyte vegetation**, **P3d Fresh-water submerged vegetation**, **P3e Fresh-water nymphaeid vegetation**, **Q51 Tall-helophyte bed**, **Q52 Small-helophyte bed**, **T15 Broadleaved swamp forest on non-acid peat**, **V14 Inundated or inundatable cropland, including rice fields**

Dominant species of EUNIS habitats: **P3b Fresh-water small pleustophyte vegetation**, **V14 Inundated or inundatable cropland, including rice fields**

Broad habitat

Occurrence in broad habitats: **Aquatic, Wetland, Forest, Synanthropic**

Distribution

Continentality amplitude: **6**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz