

Lemna trisulca

Habitus and growth type

Plant height [m]: **0.01**

Life span: **Perennial**

Life form: **Hydrophyte**

Leaf

Specific leaf area [mm²/mg]: **16.65**

Flower

Flowering period: **March-October**

Fruit, seed and dispersal

Dispersal mode: **Epizoochory**

Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Water**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.1**

Temperature indicator value: **5.5**

Moisture indicator value: **11.8**

Reaction indicator value: **6.5**

Nutrient indicator value: **5.3**

Salinity indicator value: **0.3**

Disturbance indicator values

Disturbance frequency: **1.55**

Disturbance frequency (herb layer): **1.64**

Disturbance severity: **0.26**

Disturbance severity (herb layer): **0.25**

Mowing frequency: **0.08**

Grazing pressure: **0.03**

Soil disturbance: **0.04**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [NA \(LEM\) Lemnetea](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [P3b Fresh-water small pleustophyte vegetation](#), [P3c Fresh-water large pleustophyte vegetation](#), [P3d Fresh-water submerged vegetation](#)

Constant species of EUNIS habitats: [P3b Fresh-water small pleustophyte vegetation](#), [P3c Fresh-water large pleustophyte vegetation](#), [P3d Fresh-water submerged vegetation](#), [P3e Fresh-water nymphaeid vegetation](#)

Dominant species of EUNIS habitats: [P3b Fresh-water small pleustophyte vegetation](#), [P3c Fresh-water large pleustophyte vegetation](#)

Broad habitat

Occurrence in broad habitats: **Aquatic, Wetland, Synanthropic**

Distribution

Continentality amplitude: 7