

# *Leopoldia weissii*

## Habitus and growth type

Life span: **Perennial**

Life form: **Geophyte**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Salinity relationship: **Slightly saline or brackish**

### Ellenberg-type indicator values

Light indicator value: **7.3**

Temperature indicator value: **11**

Moisture indicator value: **3**

Reaction indicator value: **8**

Nutrient indicator value: **4.7**

Salinity indicator value: **1**

### Disturbance indicator values

Disturbance frequency: **0.12**

Disturbance frequency (herb layer): **1.57**

Disturbance severity: **0.67**

Disturbance severity (herb layer): **0.22**

Mowing frequency: **0**

Grazing pressure: **0.29**

Soil disturbance: **0.16**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [FK \(PEG\) \*Pegano harmalae-Salsoletea vermiculatae\*](#)

### Broad habitat

Occurrence in broad habitats: **Scrub**