

Lepidium coronopus

Habitus and growth type

Plant height [m]: **0.13**

Life span: **Annual, Biennial or short-lived**

Life form: **Hemicryptophyte, Therophyte**

Leaf

Specific leaf area [mm²/mg]: **23.14**

Flower

Flowering period: **February-June**

Fruit, seed and dispersal

Seed mass [mg]: **3.71**

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **1, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Slightly saline or brackish**

Ellenberg-type indicator values

Light indicator value: **7.9**

Moisture indicator value: **4.6**

Reaction indicator value: **6.3**

Nutrient indicator value: **6.8**

Salinity indicator value: **0.9**

Disturbance indicator values

Disturbance frequency: **2.36**

Disturbance frequency (herb layer): **2.36**

Disturbance severity: **0.75**

Disturbance severity (herb layer): **0.75**

Mowing frequency: **0.96**

Grazing pressure: **0.23**

Soil disturbance: **0.53**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [QE \(POL\) *Polygono-Poetea annuae*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [V35 Trampled mesophilous grassland with annuals](#)

Broad habitat

Occurrence in broad habitats: **Coastal saltmarsh, Wetland, Saline vegetation, Synanthropic**

Distribution

Continentality amplitude: **5**