

# ***Limonium articulatum***

## **Habitus and growth type**

Plant height [m]: **0.14**

Life span: **Perennial**

Life form: **Chamaephyte**

## **Flower**

Flowering period: **May-August**

## **Fruit, seed and dispersal**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## **Trophic mode**

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## **Taxon origin**

Origin in Europe: **Native**

## **Ecology**

### **Environmental relationships**

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Saline**

### **Ellenberg-type indicator values**

Light indicator value: **8**

Temperature indicator value: **9**

Moisture indicator value: **5**

Reaction indicator value: **5**

Nutrient indicator value: **5**

Salinity indicator value: **6**

### **Disturbance indicator values**

Disturbance frequency: **0.69**

Disturbance frequency (herb layer): **0.74**

Disturbance severity: **0.32**

Disturbance severity (herb layer): **0.29**

Mowing frequency: **0.1**

Grazing pressure: **0.14**

Soil disturbance: **0.12**

## **Habitat and sociology**

### Syntaxon

Diagnostic species of phytosociological classes: [\*\*JB \(CRI\) \*Crithmo-Staticetea\*\*\*](#)

### Broad habitat

Occurrence in broad habitats: **Coastal saltmarsh, Coastal beach, dune or shingle, Coastal cliff, Saline vegetation, Sparsely vegetated (incl. rock and scree)**