

Limonium dodartii

Habitus and growth type

Life span: **Perennial**

Life form: **Hemicryptophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Saline**

Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **6**

Moisture indicator value: **7**

Reaction indicator value: **7**

Nutrient indicator value: **5**

Salinity indicator value: **7**

Disturbance indicator values

Disturbance frequency: **1.33**

Disturbance frequency (herb layer): **1.33**

Disturbance severity: **0.46**

Disturbance severity (herb layer): **0.46**

Mowing frequency: **0**

Grazing pressure: **0.3**

Soil disturbance: **0.23**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [JB \(CRI\) Crithmo-Staticetea](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [N31 Atlantic and Baltic rocky sea cliff and shore](#)

Broad habitat

Occurrence in broad habitats: **Coastal cliff**