

Limonium suffruticosum

Habitus and growth type

Life span: **Perennial**

Life form: **Chamaephyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Ellenberg-type indicator values

Light indicator value: **8.5**

Temperature indicator value: **6**

Moisture indicator value: **2.3**

Disturbance indicator values

Disturbance frequency: **2.31**

Disturbance frequency (herb layer): **2.29**

Disturbance severity: **0.42**

Disturbance severity (herb layer): **0.42**

Mowing frequency: **0.07**

Grazing pressure: **0.55**

Soil disturbance: **0.3**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [EC \(KAL\) *Kalidietea foliati*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R64 Semi-desert salt pan](#)

Constant species of EUNIS habitats: [R64 Semi-desert salt pan](#)

Broad habitat

Occurrence in broad habitats: **Saline vegetation**