

# *Limonium vulgare*

## Habitus and growth type

Plant height [m]: **0.17**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **10.71**

## Fruit, seed and dispersal

Seed mass [mg]: **1.18**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Wet**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Saline**

### Ellenberg-type indicator values

Light indicator value: **8.7**

Temperature indicator value: **5.5**

Moisture indicator value: **7.7**

Reaction indicator value: **7.7**

Nutrient indicator value: **5.7**

Salinity indicator value: **7.3**

### Disturbance indicator values

Disturbance frequency: **2.15**

Disturbance frequency (herb layer): **2.21**

Disturbance severity: **0.5**

Disturbance severity (herb layer): **0.5**

Mowing frequency: **0.01**

Grazing pressure: **0.3**

Soil disturbance: **0.2**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [MF \(JUN\) \*Juncetea maritimi\*](#)

### EUNIS habitat

Diagnostic species of EUNIS habitats: [MA224 Atlantic mid-low saltmarsh](#), [MA253 Mediterranean mid-low saltmarsh](#)

Constant species of EUNIS habitats: [MA221 Atlantic saltmarsh driftline](#), [MA223 Atlantic upper-mid saltmarsh and saline and brackish reed, rush and sedge bed](#), [MA224 Atlantic mid-low saltmarsh](#), [MA252 Mediterranean upper-mid saltmarsh and saline and brackish reed, rush and sedge bed](#), [MA253 Mediterranean mid-low saltmarsh](#)

### Broad habitat

Occurrence in broad habitats: **Coastal saltmarsh**

## Distribution

Continentality amplitude: **6**