

# *Linaria biebersteinii*

## Habitus and growth type

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Ellenberg-type indicator values

Light indicator value: **8.3**

Temperature indicator value: **5.1**

Moisture indicator value: **5.1**

Disturbance indicator values

Disturbance frequency: **1.57**

Disturbance frequency (herb layer): **2.01**

Disturbance severity: **0.57**

Disturbance severity (herb layer): **0.56**

Mowing frequency: **0.79**

Grazing pressure: **0.34**

Soil disturbance: **0.25**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [\*\*DA \(FES\) Festuco-Brometea\*\*](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**