Linaria propinqua

Habitus and growth type

Life span: **Annual**
Life form: **Therophyte**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Ellenberg-type indicator values

Light indicator value: **9**
Temperature indicator value: **5**
Moisture indicator value: **1**
Reaction indicator value: **7**
Nutrient indicator value: **1**

Disturbance indicator values

Disturbance frequency: **1.2**
Disturbance frequency (herb layer): **1.73**
Disturbance severity: **0.5**
Disturbance severity (herb layer): **0.47**
Mowing frequency: **0.5**
Grazing pressure: **0.25**
Soil disturbance: **0.2**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **KE (THL) *Thlaspietea rotundifolii***

Broad habitat

Occurrence in broad habitats: **Sparsely vegetated (incl. rock and scree)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz