# *Linaria spartea*

### Habitus and growth type

Plant height [m]: **0.38**
Life span: **Annual**
Life form: **Therophyte**

### Fruit, seed and dispersal

Seed mass [mg]: **0.04**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

### Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

### Taxon origin

Origin in Europe: **Native**

### Ecology

#### Ellenberg-type indicator values

Light indicator value: **9**
Temperature indicator value: **7**
Moisture indicator value: **1**
Reaction indicator value: **4**
Nutrient indicator value: **4**
Salinity indicator value: **0**

#### Disturbance indicator values

Disturbance frequency: **0.99**
Disturbance frequency (herb layer): **1.85**
Disturbance severity: **0.44**
Disturbance severity (herb layer): **0.38**
Mowing frequency: **0.55**
Grazing pressure: **0.29**
Soil disturbance: **0.24**

### Habitat and sociology

#### Syntaxon

Diagnostic species of phytosociological classes: **FI (TUB) *Helianthemetea guttati***

#### EUNIS habitat

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*

Diagnostic species of EUNIS habitats: **R1R Mediterranean to Atlantic open, dry, acid and neutral grassland**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**

## Distribution

Continentality: **2**

Continentality amplitude: **2**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*