

Malva sylvestris

Habitus and growth type

Plant height [m]: **0.76**

Life span: **Biennial or short-lived, Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **22.61**

Flower

Flowering period: **April-August**

Fruit, seed and dispersal

Seed mass [mg]: **5.29**

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **2, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.1**

Temperature indicator value: **6.2**

Reaction indicator value: **7.1**

Nutrient indicator value: **7.6**

Salinity indicator value: **0.1**

Disturbance indicator values

Disturbance frequency: **1.41**

Disturbance frequency (herb layer): **1.89**

Disturbance severity: **0.81**

Disturbance severity (herb layer): **0.79**

Mowing frequency: **0.89**

Grazing pressure: **0.17**

Soil disturbance: **0.5**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [QB \(SIS\) *Sisymbrietea*](#), [QC \(CHE\) *Chenopodietea*](#), [QF \(ART\) *Artemisietea vulgaris*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [V33 Dry Mediterranean land with unpalatable non-vernal herbaceous vegetation](#)

Broad habitat

Occurrence in broad habitats: **Scrub, Synanthropic**

Distribution

Continentality amplitude: **6**