

Marrubium vulgare

Habitus and growth type

Plant height [m]: **0.4**

Life span: **Perennial**

Life form: **Chamaephyte, Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **14.2**

Flower

Flowering period: **April-August**

Fruit, seed and dispersal

Seed mass [mg]: **0.93**

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **2, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.7**

Moisture indicator value: **3.8**

Reaction indicator value: **7.8**

Nutrient indicator value: **7.8**

Salinity indicator value: **0.1**

Disturbance indicator values

Disturbance frequency: **1.17**

Disturbance frequency (herb layer): **1.76**

Disturbance severity: **0.76**

Disturbance severity (herb layer): **0.71**

Mowing frequency: **0.53**

Grazing pressure: **0.21**

Soil disturbance: **0.38**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [QF \(ART\) Artemisietea vulgaris](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree), Synanthropic**

Distribution

Continentality amplitude: **6**