

Medicago scutellata

Habitus and growth type

Plant height [m]: **0.09**

Life span: **Annual**

Life form: **Therophyte**

Flower

Flowering period: **March-May**

Fruit, seed and dispersal

Seed mass [mg]: **18.5**

Dispersal mode: **Epizoochory**

Dispersal distance class: **6**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.7**

Moisture indicator value: **5**

Reaction indicator value: **6.7**

Nutrient indicator value: **6**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.56**

Disturbance frequency (herb layer): **2.17**

Disturbance severity: **0.86**

Disturbance severity (herb layer): **0.84**

Mowing frequency: **0.64**

Grazing pressure: **0.17**

Soil disturbance: **0.58**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [QC \(CHE\) Chenopodietea](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Sparsely vegetated (incl. rock and scree), Synanthropic**