Melampyrum velebiticum

Habitus and growth type

Plant height [m]: **0.3**
Life span: **Annual**
Life form: **Therophyte**

Flower

Flowering period: **May-August**

Fruit, seed and dispersal

Dispersal mode: **Myrmecochory**
Dispersal distance class: **3**

Trophic mode

Parasitism and mycoheterotrophy: **euphytoid hemiparasite**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **6**
Temperature indicator value: **6.5**
Moisture indicator value: **6**
Reaction indicator value: **7**
Nutrient indicator value: **5**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.03**
Disturbance frequency (herb layer): **1.31**
Disturbance severity: **0.69**
Disturbance severity (herb layer): **0.11**
Mowing frequency: **0.01**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Grazing pressure: **0.19**
Soil disturbance: **0.1**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **CL (GER) *Trifolio-Geranietea sanguinei***

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Forest**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz