Minuartia glomerata

Habitus and growth type

Life span: **Annual, Biennial or short-lived, Perennial**
Life form: **Hemicryptophyte, Therophyte**

Fruit, seed and dispersal

Seed mass [mg]: **0.08**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.8**
Temperature indicator value: **5.8**
Moisture indicator value: **3.6**
Reaction indicator value: **6.5**
Nutrient indicator value: **2.5**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.75**
Disturbance frequency (herb layer): **2.14**
Disturbance severity: **0.43**
Disturbance severity (herb layer): **0.42**
Mowing frequency: **0.3**
Grazing pressure: **0.39**
Soil disturbance: **0.23**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Habitat and sociology

EUNIS habitat

Diagnostic species of EUNIS habitats: **R11 Pannonian and Pontic sandy steppe**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz