

# *Moehringia pentandra*

## Habitus and growth type

Plant height [m]: **0.15**

Life span: **Annual**

Life form: **Therophyte**

## Flower

Flowering period: **March-July**

## Fruit, seed and dispersal

Dispersal mode: **Myrmecochory**

Dispersal distance class: **3**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **4**

Temperature indicator value: **8**

Moisture indicator value: **5.5**

Reaction indicator value: **5**

Nutrient indicator value: **7**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.05**

Disturbance frequency (herb layer): **1.32**

Disturbance severity: **0.68**

Disturbance severity (herb layer): **0.12**

Mowing frequency: **0.01**

Grazing pressure: **0.25**

Soil disturbance: **0.1**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [\*\*CA \(FAG\) \*Carpino-Fagetetea sylvaticae\*\*\*](#),  
[\*\*FA \(QUI\) \*Quercetea ilicis\*\*\*](#)

### Broad habitat

Occurrence in broad habitats: **Scrub, Forest, Sparsely vegetated (incl. rock and scree)**