

Moenchia erecta

Habitus and growth type

Plant height [m]: **0.09**

Life span: **Annual**

Life form: **Therophyte**

Flower

Flowering period: **April-June**

Fruit, seed and dispersal

Seed mass [mg]: **0.08**

Dispersal mode: **Myrmecochory**

Dispersal distance class: **3**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Acidic**

Nutrient relationship: **Dystrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **9**

Temperature indicator value: **7.6**

Moisture indicator value: **2.2**

Reaction indicator value: **3.6**

Nutrient indicator value: **1.6**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.79**

Disturbance frequency (herb layer): **2.16**

Disturbance severity: **0.46**

Disturbance severity (herb layer): **0.45**

Mowing frequency: **0.47**

Grazing pressure: **0.39**

Soil disturbance: **0.32**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [**CJ \(COR\) Koelerio-Corynephoretea**](#)
[**canescens**](#), [**FI \(TUB\) Helianthemetea guttati**](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [**R1R Mediterranean to Atlantic open, dry, acid and neutral grassland**](#)

Constant species of EUNIS habitats: [**R1R Mediterranean to Atlantic open, dry, acid and neutral grassland**](#)

Broad habitat

Occurrence in broad habitats: **Wetland, Grassland (non-alpine, non-saline)**

Distribution

Continentality: **3**

Continentality amplitude: **3**