

# *Molineriella laevis*

## Habitus and growth type

Life span: **Annual**

Life form: **Therophyte**

## Fruit, seed and dispersal

Seed mass [mg]: **0.03**

Dispersal mode: **Local non-specific dispersal**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

Ellenberg-type indicator values

Light indicator value: **9**

Temperature indicator value: **7**

Moisture indicator value: **1**

Reaction indicator value: **3**

Nutrient indicator value: **2**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **2.09**

Disturbance frequency (herb layer): **2.15**

Disturbance severity: **0.38**

Disturbance severity (herb layer): **0.38**

Mowing frequency: **0.54**

Grazing pressure: **0.34**

Soil disturbance: **0.26**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [FI \(TUB\) \*Helianthemetea guttati\*](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [R1N Open Iberian supramediterranean dry acid and neutral grassland](#), [R1R Mediterranean to Atlantic open, dry, acid and](#)

**neutral grassland**

Constant species of EUNIS habitats: **R1N Open Iberian supramediterranean dry acid and neutral grassland**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**