

# *Morina persica*

## Habitus and growth type

Life span: **Perennial**

Life form: **Hemicryptophyte**

## Fruit, seed and dispersal

Seed mass [mg]: **17.15**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Ellenberg-type indicator values

Light indicator value: **8.2**

Temperature indicator value: **7**

Moisture indicator value: **3.7**

Reaction indicator value: **6.8**

Nutrient indicator value: **3.2**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.89**

Disturbance frequency (herb layer): **1.87**

Disturbance severity: **0.46**

Disturbance severity (herb layer): **0.38**

Mowing frequency: **0.1**

Grazing pressure: **0.33**

Soil disturbance: **0.15**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [DA \(FES\) Festuco-Brometea](#), [FR \(DAP\) Daphno-Festucetea](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline),  
Oromediterranean scrub**