Muscari neglectum

Habitus and growth type

Plant height [m]: **0.16**
Life span: **Perennial**
Life form: **Geophyte**
Spinescence: **not spinescent**

Leaf

Specific leaf area [mm^2/mg]: **12.54**

Flower

Flowering period: **February-June**

Fruit, seed and dispersal

Seed mass [mg]: **3.17**
Dispersal mode: **Local non-specific dispersal, Anthropochory**
Dispersal distance class: **1, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.3**
Temperature indicator value: **7.1**
Moisture indicator value: **3.4**
Reaction indicator value: **7**
Salinity indicator value: **0.2**

Disturbance indicator values

Disturbance frequency: **0.86**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Disturbance frequency (herb layer): **1.75**
Disturbance severity: **0.52**
Disturbance severity (herb layer): **0.42**
Mowing frequency: **0.27**
Grazing pressure: **0.33**
Soil disturbance: **0.21**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **DA (FES) *Festuco-Brometea***, **FP (ONO) *Festuco hystricis-Ononidetea striatae***

EUNIS habitat

Diagnostic species of EUNIS habitats: **R14 Perennial rocky grassland of the Italian Peninsula**
Constant species of EUNIS habitats: **R14 Perennial rocky grassland of the Italian Peninsula**

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree)**

Distribution

Continentality amplitude: **8**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz