# *Myosotis secunda*

## Habitus and growth type

Plant height [m]: **0.3**
Life span: **Perennial**
Life form: **Hemicryptophyte**

## Fruit, seed and dispersal

Seed mass [mg]: **0.2**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Wet**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7.7**
Temperature indicator value: **6**
Moisture indicator value: **9.3**
Reaction indicator value: **4.7**
Nutrient indicator value: **3.7**
Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.14**
Disturbance frequency (herb layer): **1.73**
Disturbance severity: **0.42**
Disturbance severity (herb layer): **0.38**
Mowing frequency: **0.37**
Grazing pressure: **0.24**
Soil disturbance: **0.17**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: **OB (LIT) *Littorelletea uniflorae***

### Broad habitat

Occurrence in broad habitats: **Aquatic**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*