Myosotis sparsiflora

Habitus and growth type

Plant height [m]: **0.29**
Life span: **Annual**
Life form: **Therophyte**

Fruit, seed and dispersal

Seed mass [mg]: **0.47**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **5.4**
Temperature indicator value: **5.8**
Moisture indicator value: **6**
Reaction indicator value: **7.3**
Nutrient indicator value: **7**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.23**
Disturbance frequency (herb layer): **1.36**
Disturbance severity: **0.66**
Disturbance severity (herb layer): **0.29**
Mowing frequency: **0.12**
Grazing pressure: **0.19**
Soil disturbance: **0.15**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **HA (POP) *Alno glutinosae-Populetea albae***

Broad habitat

Occurrence in broad habitats: **Forest, Synanthropic**

Distribution

Continentality: **7**
Continentality amplitude: **3**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz