

Myosotis sylvatica

Habitus and growth type

Plant height [m]: **0.32**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **29.68**

Flower

Flowering period: **March-September**

Fruit, seed and dispersal

Seed mass [mg]: **0.48**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Mesic**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **5.5**

Temperature indicator value: **4.6**

Moisture indicator value: **5.6**

Reaction indicator value: **6.2**

Nutrient indicator value: **6.7**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.16**

Disturbance frequency (herb layer): **1.28**

Disturbance severity: **0.63**

Disturbance severity (herb layer): **0.21**

Mowing frequency: **0.08**

Grazing pressure: **0.19**

Soil disturbance: **0.13**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [CA \(FAG\) *Carpino-Fagetea sylvaticae*](#),
[DA \(FES\) *Festuco-Brometea*](#), [QG \(EPI\) *Epilobietea angustifolii*](#)

EUNIS habitat

Constant species of EUNIS habitats: [R56 Montane to subalpine moist or wet tall-herb and fern fringe](#), [S25 Subalpine and subarctic deciduous scrub](#), [T33 Mediterranean mountain Abies forest](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Alpine-subalpine and arctic grassland, Scrub, Forest, Sparsely vegetated (incl. rock and scree), Synanthropic**

Distribution

Continentality: **5**

Continentality amplitude: **3**