Myriolimon ferulaceum

Habitus and growth type

Plant height [m]: **0.25**
Life span: **Perennial**
Life form: **Chamaephyte**

Flower

Flowering period: **May-July**

Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **native**

Ecology

Environmental relationships

Substrate humidity relationship: **Wet**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Saline**

Ellenberg-type indicator values

Light indicator value: **8.5**
Temperature indicator value: **10**
Moisture indicator value: **7.5**
Reaction indicator value: **8.5**
Nutrient indicator value: **6.5**
Salinity indicator value: **8**

Disturbance indicator values

Disturbance frequency: **2.3**
Disturbance frequency (herb layer): **2.3**
Disturbance severity: **0.44**
Disturbance severity (herb layer): **0.44**
Mowing frequency: **0**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Grazing pressure: **0.19**
Soil disturbance: **0.2**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **MG (SAL) *Salicornietea fruticosae***

Broad habitat

Occurrence in broad habitats: **Coastal saltmarsh, Saline vegetation**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz