

# ***Nasturtium microphyllum***

## **Habitus and growth type**

Plant height [m]: **0.33**

Life span: **Perennial**

Life form: **Hemicryptophyte**

## **Leaf**

Specific leaf area [mm<sup>2</sup>/mg]: **72.1**

## **Flower**

Flowering period: **May-August**

## **Fruit, seed and dispersal**

Seed mass [mg]: **0.13**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## **Trophic mode**

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## **Taxon origin**

Origin in Europe: **native**

## **Ecology**

### **Environmental relationships**

Substrate humidity relationship: **Water**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Eutrophic**

Salinity relationship: **Non-saline**

### **Ellenberg-type indicator values**

Light indicator value: **7.3**

Temperature indicator value: **5**

Moisture indicator value: **10**

Reaction indicator value: **6.5**

Nutrient indicator value: **5.5**

Salinity indicator value: **0.3**

### **Disturbance indicator values**

Disturbance frequency: **1.58**

Disturbance frequency (herb layer): **1.75**

Disturbance severity: **0.29**

Disturbance severity (herb layer): **0.28**

Mowing frequency: **0.2**

Grazing pressure: **0.14**

Soil disturbance: **0.09**

## **Habitat and sociology**

### **Syntaxon**

Diagnostic species of phytosociological classes: [OD \(PHR\) \*Phragmito-Magnocaricetea\*](#)

### **Broad habitat**

Occurrence in broad habitats: **Aquatic, Wetland, Sparsely vegetated (incl. rock and scree)**

## **Distribution**

Continentality: **3**

Continentality amplitude: **3**