# *Oenanthe pimpinelloides*

## Habitus and growth type

Plant height [m]: **0.57**
Life span: **Perennial**
Life form: **Hemicryptophyte**

## Flower

Flowering period: **April-July**

## Fruit, seed and dispersal

Seed mass [mg]: **2**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Mesic**
Substrate reaction relationship: **Slightly acidic to near-neutral**
Nutrient relationship: **Mesotrophic**
Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **6.7**
Temperature indicator value: **8.3**
Moisture indicator value: **6**
Reaction indicator value: **5.5**
Nutrient indicator value: **4**
Salinity indicator value: **0.3**

### Disturbance indicator values

Disturbance frequency: **0.18**
Disturbance frequency (herb layer): **1.39**
Disturbance severity: **0.65**
Disturbance severity (herb layer): **0.24**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*

Mowing frequency: **0.1**
Grazing pressure: **0.23**
Soil disturbance: **0.15**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: **CM (MOL) *Molinio-Arrhenatheretea***

### EUNIS habitat

Diagnostic species of EUNIS habitats: **S34 Balkan-Anatolian submontane genistoid scrub**, **T1G Alnus cordata forest**

### Broad habitat

Occurrence in broad habitats: **Spring, Grassland (non-alpine, non-saline), Scrub, Forest**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*