

Oenothera glazioviana

Habitus and growth type

Plant height [m]: **0.98**

Life span: **Biennial or short-lived**

Life form: **Hemicryptophyte**

Leaf

Specific leaf area [mm²/mg]: **24.21**

Flower

Flowering period: **May-September**

Fruit, seed and dispersal

Seed mass [mg]: **0.62**

Dispersal mode: **Local non-specific dispersal, Anthropochory**

Dispersal distance class: **2, 7**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Neophyte**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Slightly acidic to near-neutral**

Nutrient relationship: **Mesotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.8**

Temperature indicator value: **6.8**

Moisture indicator value: **3.5**

Reaction indicator value: **6.3**

Nutrient indicator value: **4.8**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.12**

Disturbance frequency (herb layer): **1.67**

Disturbance severity: **0.65**

Disturbance severity (herb layer): **0.61**

Mowing frequency: **0.55**

Grazing pressure: **0.19**

Soil disturbance: **0.3**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [QF \(ART\) Artemisietea vulgaris](#)

Broad habitat

Occurrence in broad habitats: **Scrub, Sparsely vegetated (incl. rock and scree),
Synanthropic**