

# *Ononis fruticosa*

## Habitus and growth type

Plant height [m]: **0.65**

Life span: **Perennial**

Life form: **Phanerophyte, Shrub**

Spinescence: **not spinescent**

## Leaf

Specific leaf area [mm<sup>2</sup>/mg]: **12.88**

## Flower

Flowering period: **April-August**

## Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **2**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

## Taxon origin

Origin in Europe: **native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **8**

Temperature indicator value: **7.5**

Moisture indicator value: **3**

Reaction indicator value: **8**

Nutrient indicator value: **3**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.53**

Disturbance frequency (herb layer): **1.46**

Disturbance severity: **0.62**

Disturbance severity (herb layer): **0.22**

Mowing frequency: **0.05**

Grazing pressure: **0.36**

Soil disturbance: **0.16**

## **Habitat and sociology**

### Syntaxon

Diagnostic species of phytosociological classes: [\*\*CB \(PUB\) Quercetea pubescentis\*\*](#), [\*\*FB \(ROS\) Ononido-Rosmarinetea\*\*](#)

### Broad habitat

Occurrence in broad habitats: **Scrub, Forest**