

Ononis pusilla

Habitus and growth type

Plant height [m]: **0.21**

Life span: **Perennial**

Life form: **Hemicryptophyte**

Spinescence: **not spinescent**

Leaf

Specific leaf area [mm²/mg]: **13.26**

Flower

Flowering period: **May-September**

Fruit, seed and dispersal

Seed mass [mg]: **3**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **8.5**

Temperature indicator value: **7.4**

Moisture indicator value: **2.9**

Reaction indicator value: **8.3**

Nutrient indicator value: **1.5**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.81**

Disturbance frequency (herb layer): **1.67**

Disturbance severity: **0.53**

Disturbance severity (herb layer): **0.4**

Mowing frequency: **0.37**

Grazing pressure: **0.37**

Soil disturbance: **0.19**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [**DA \(FES\) Festuco-Brometea**](#), [**FP \(ONO\) Festuco hystricis-Ononidetea striatae**](#)

EUNIS habitat

Constant species of EUNIS habitats: [**R18 Perennial rocky calcareous grassland of subatlantic-submediterranean Europe**](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree)**