

# *Ononis verae*

## Habitus and growth type

Life span: **Annual**

Life form: **Therophyte**

## Fruit, seed and dispersal

Dispersal mode: **Local non-specific dispersal**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **symbiosis with rhizobia**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7**

Temperature indicator value: **11**

Moisture indicator value: **3**

Reaction indicator value: **8**

Nutrient indicator value: **5**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **0.53**

Disturbance frequency (herb layer): **1.62**

Disturbance severity: **0.77**

Disturbance severity (herb layer): **0.46**

Mowing frequency: **0.17**

Grazing pressure: **0.27**

Soil disturbance: **0.25**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: [EJ \(TRA\) \*Stipo-Trachynietea distachyae\*](#)

## Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline)**