

# *Onosma fastigiata*

## Habitus and growth type

Plant height [m]: **0.25**

Life span: **Biennial or short-lived**

Life form: **Hemicryptophyte**

## Flower

Flowering period: **March-July**

## Fruit, seed and dispersal

Seed mass [mg]: **9.66**

Dispersal mode: **Local non-specific dispersal**

Dispersal distance class: **1**

## Trophic mode

Parasitism and mycoheterotrophy: **autotroph**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **Native**

## Ecology

### Environmental relationships

Substrate humidity relationship: **Dry**

Substrate reaction relationship: **Alkaline**

Nutrient relationship: **Oligotrophic**

Salinity relationship: **Non-saline**

### Ellenberg-type indicator values

Light indicator value: **7**

Temperature indicator value: **6.5**

Moisture indicator value: **3**

Reaction indicator value: **8**

Nutrient indicator value: **2**

Salinity indicator value: **0**

### Disturbance indicator values

Disturbance frequency: **1.05**

Disturbance frequency (herb layer): **1.74**

Disturbance severity: **0.51**

Disturbance severity (herb layer): **0.46**

Mowing frequency: **0.66**

Grazing pressure: **0.39**

Soil disturbance: **0.22**

## **Habitat and sociology**

### Syntaxon

Diagnostic species of phytosociological classes: [DA \(FES\) Festuco-Brometea](#)

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Sparsely vegetated (incl. rock and scree)**