# *Ornithogalum comosum*

### Habitus and growth type

Plant height [m]: **0.2**
Life span: **Perennial**
Life form: **Geophyte**

### Flower

Flowering period: **March-June**

### Fruit, seed and dispersal

Seed mass [mg]: **3.51**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **1**

### Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

### Taxon origin

Origin in Europe: **Native**

### Ecology

#### Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Oligotrophic**
Salinity relationship: **Non-saline**

#### Ellenberg-type indicator values

Light indicator value: **8.3**
Temperature indicator value: **8.5**
Moisture indicator value: **2**
Reaction indicator value: **8**
Nutrient indicator value: **1.5**
Salinity indicator value: **0**

#### Disturbance indicator values

Disturbance frequency: **1.82**
Disturbance frequency (herb layer): **2.23**
Disturbance severity: **0.54**
Disturbance severity (herb layer): **0.53**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*

Mowing frequency: **0.41**
Grazing pressure: **0.34**
Soil disturbance: **0.29**

## Habitat and sociology

### Syntaxon

Diagnostic species of phytosociological classes: **DA (FES) *Festuco-Brometea***

### EUNIS habitat

Diagnostic species of EUNIS habitats: **R19 Dry steppic submediterranean pasture of the Amphi-Adriatic region**

### Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub, Sparsely vegetated (incl. rock and scree)**

*© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz*