Ornithogalum narbonense

Habitus and growth type

Plant height [m]: **0.5**
Life span: **Perennial**
Life form: **Geophyte**

Flower

Flowering period: **April-June**

Fruit, seed and dispersal

Seed mass [mg]: **3.22**
Dispersal mode: **Local non-specific dispersal**
Dispersal distance class: **2**

Trophic mode

Parasitism and mycoheterotrophy: **autotroph**
Carnivory: **non-carnivorous**
Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate humidity relationship: **Dry**
Substrate reaction relationship: **Alkaline**
Nutrient relationship: **Eutrophic**
Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.3**
Temperature indicator value: **8.5**
Moisture indicator value: **4.3**
Reaction indicator value: **7.3**
Nutrient indicator value: **5.9**
Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.08**
Disturbance frequency (herb layer): **1.87**
Disturbance severity: **0.8**
Disturbance severity (herb layer): **0.73**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz

Mowing frequency: **0.43**
Grazing pressure: **0.25**
Soil disturbance: **0.42**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: **DA (FES) *Festuco-Brometea***, **FF (LYG) *Lygeo sparti-Stipetea tenacissimae***

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline), Scrub**

© 2014–2026 Pladias – Citation: Pladias – Database of the Czech Flora and Vegetation. www.pladias.cz