

# *Orobanche latisquama*

## Habitus and growth type

Life span: **Perennial**

Life form: **Geophyte**

## Fruit, seed and dispersal

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

## Trophic mode

Parasitism and mycoheterotrophy: **obligate root parasite**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

## Taxon origin

Origin in Europe: **native**

## Ecology

Ellenberg-type indicator values

Light indicator value: **8.3**

Temperature indicator value: **8.3**

Moisture indicator value: **2.3**

Reaction indicator value: **7**

Nutrient indicator value: **2.9**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **0.51**

Disturbance frequency (herb layer): **1.72**

Disturbance severity: **0.7**

Disturbance severity (herb layer): **0.19**

Mowing frequency: **0**

Grazing pressure: **0.39**

Soil disturbance: **0.16**

## Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [FB \(ROS\) Ononido-Rosmarinetea](#)

EUNIS habitat

Diagnostic species of EUNIS habitats: [S61 Western basiphilous garrigue](#)

Broad habitat

Occurrence in broad habitats: **Scrub**