

Orobanche reticulata

Habitus and growth type

Plant height [m]: **0.36**

Life span: **Perennial**

Life form: **Geophyte**

Flower

Flowering period: **April-July**

Fruit, seed and dispersal

Dispersal mode: **Anemochory**

Dispersal distance class: **5**

Trophic mode

Parasitism and mycoheterotrophy: **obligate root parasite**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in Europe: **Native**

Ecology

Environmental relationships

Substrate reaction relationship: **Alkaline**

Salinity relationship: **Non-saline**

Ellenberg-type indicator values

Light indicator value: **7.3**

Reaction indicator value: **7.4**

Salinity indicator value: **0**

Disturbance indicator values

Disturbance frequency: **1.04**

Disturbance frequency (herb layer): **1.61**

Disturbance severity: **0.45**

Disturbance severity (herb layer): **0.4**

Mowing frequency: **0.24**

Grazing pressure: **0.25**

Soil disturbance: **0.17**

Habitat and sociology

Syntaxon

Diagnostic species of phytosociological classes: [DA \(FES\) Festuco-Brometea](#)

Broad habitat

Occurrence in broad habitats: **Grassland (non-alpine, non-saline),
Oromediterranean scrub, Scrub**

Distribution

Continentality: **6**

Continentality amplitude: **6**